

Eigenvalue problems of Nordhaus-Gaddum type

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis,
Memphis TN 38152, USA

February 1, 2008

Abstract

Let G be a graph with n vertices and m edges and let $\mu_1(G) \geq \dots \geq \mu_n(G)$ be the eigenvalues of its adjacency matrix. We discuss the following general problem. For k fixed and n large, find or estimate

$$f_k(n) = \max_{v(G)=n} |\mu_k(G)| + |\mu_k(\overline{G})|.$$

In particular we prove that

$$\frac{4}{3}n - 2 \leq f_1(n) < (\sqrt{2} - c)n$$

for some $c > 8 \times 10^{-7}$ independent of n . We also show that

$$\begin{aligned} \frac{\sqrt{2}}{2}n - 3 &< f_2(n) < \frac{\sqrt{2}}{2}n, \\ \frac{\sqrt{2}}{2}n - 3 &< f_n(n) \leq \frac{\sqrt{3}}{2}n. \end{aligned}$$

AMS classification: 15A42, 05C50

Keywords: graph eigenvalues, complementary graph, maximum eigenvalue, minimum eigenvalue, Nordhaus-Gaddum problems,

1 Introduction

Our notation is standard (e.g., see [1], [2], and [3]); in particular, all graphs are defined on the vertex set $\{1, 2, \dots, n\} = [n]$ and $G(n, m)$ stands for a graph with n vertices and m edges. We write $\Gamma(u)$ for the set of neighbors of the vertex u and set $d(u) = |\Gamma(u)|$. Given a graph G of order n , we assume that the eigenvalues of the adjacency matrix of G are ordered as $\mu(G) = \mu_1(G) \geq \dots \geq \mu_n(G)$. As usual, $\overline{G}$ denotes the complement of a graph G and $\omega(G)$ stands for the clique number of G .

Nosal [10] showed that for every graph G of order n ,

$$n - 1 \leq \mu(G) + \mu(\overline{G}) < \sqrt{2}n. \quad (1)$$

Quite of attention has been given to second of these inequalities. In [8] it was shown that

$$\mu(G) + \mu(\overline{G}) \leq \sqrt{\left(2 - \frac{1}{\omega(G)} - \frac{1}{\omega(\overline{G})}\right) n(n-1)}, \quad (2)$$

improving earlier results in [4], [5], [7], and [11]. Unfortunately inequality (2) is not much better than (1) when both $\omega(G)$ and $\omega(\overline{G})$ are large enough. Thus, it is natural to ask whether $\sqrt{2}$ in (1) can be replaced by a smaller absolute constant for n sufficiently large. In this note we answer this question in the positive but first we state a more general problem.

Problem 1 *For every $1 \leq k \leq n$ find*

$$f_k(n) = \max_{v(G)=n} |\mu_k(G)| + |\mu_k(\overline{G})|.$$

It is difficult to determine precisely $f_k(n)$ for every n and k , so at this stage it seems more practical to estimate it asymptotically. In this note we show that

$$\frac{4}{3}n - 2 \leq f_1(n) < (\sqrt{2} - c)n \quad (3)$$

for some $c > 8 \times 10^{-7}$ independent of n . For $f_2(n)$ we give the following tight bounds

$$\frac{\sqrt{2}}{2}n - 3 < f_2(n) < \frac{\sqrt{2}}{2}n. \quad (4)$$

We also show that

$$\frac{\sqrt{2}}{2}n - 3 < f_n(n) \leq \frac{\sqrt{3}}{2}n \quad (5)$$

Finally for fixed k , $2 < k < n$, and n large, we prove that

$$\begin{aligned} \left\lfloor \frac{n}{k} \right\rfloor - 1 &\leq f_k(n) \leq \sqrt{\frac{2}{k}}n, \\ \left\lfloor \frac{n}{k} \right\rfloor + 1 &\leq f_{n-k}(n) \leq \sqrt{\frac{2}{k}}n. \end{aligned}$$

2 Bounds on $f_1(n)$

Before stating the main result of this section, we shall recall two auxiliary results whose proofs can be found in [9]. Given a graph $G = G(n, m)$, let

$$s(G) = \sum_{u \in V(G)} \left| d(u) - \frac{2m}{n} \right|.$$

Proposition 2 For every graph $G = G(n, m)$,

$$\frac{s^2(G)}{2n^2\sqrt{2m}} \leq \mu_1(G) - \frac{2m}{n} \leq \sqrt{s(G)}, \quad (6)$$

and

$$\mu_n(G) + \mu_n(\overline{G}) \leq -1 - \frac{s^2(G)}{n^3}. \quad (7)$$

Decreasing the constant $\sqrt{2}$ in (1) happened to be a surprisingly challenging task for the author. The little progress that has been made is given in the following theorem.

Theorem 3 There exists $c \geq 8 \times 10^{-7}$ such that

$$\mu_1(G) + \mu_1(\overline{G}) \leq (\sqrt{2} - c)n.$$

for every graph G of order n .

Proof Assume the opposite: let $\varepsilon = 8 \times 10^{-7}$ and let there exist a graph G of order n such that

$$\mu_1(G) + \mu_1(\overline{G}) > (\sqrt{2} - \varepsilon)n.$$

Writing $A(G)$ for the adjacency matrix of G , we have

$$\sum_{i=1}^n \mu_i^2(G) = \text{tr}(A^2(G)) = 2e(G), \quad (8)$$

implying that

$$\mu_1^2(G) + \mu_n^2(G) + \mu_1^2(\overline{G}) + \mu_n^2(\overline{G}) \leq 2e(G) + 2e(\overline{G}) < n^2.$$

From

$$\mu_1^2(G) + \mu_1^2(\overline{G}) \geq \frac{1}{2}(\mu_1(G) + \mu_1(\overline{G}))^2 > \left(1 - \frac{\varepsilon}{\sqrt{2}}\right)^2 n^2 > (1 - 2\varepsilon)n^2$$

we find that

$$|\mu_n(G)| + |\mu_n(\overline{G})| \leq \sqrt{2(\mu_n^2(G) + \mu_n^2(\overline{G}))} < \sqrt{4\varepsilon}n, \quad (9)$$

and so, $\mu_n(G) + \mu_n(\overline{G}) > -\sqrt{4\varepsilon}n$. We thus have $\sqrt{4\varepsilon}n^4 \geq s^2(G)$. On the other hand, by (6) and in view of $s(G) = s(\overline{G})$, we see that

$$\mu_1(G) + \mu_1(\overline{G}) \leq n - 1 + 2\sqrt{s(G)} < n + 2\sqrt{s(G)},$$

and, by (9), it follows that

$$(\sqrt{2} - \varepsilon)n < n + 2(4\varepsilon)^{1/8}n.$$

Dividing by n , we obtain $(\sqrt{2} - 1) < \varepsilon + 2^{5/4}\varepsilon^{1/8}$, a contradiction for $\varepsilon = 8 \times 10^{-7}$. $\square$

It is certain that the upper bound given by Theorem 3 is far from the best one. We shall give below a lower bound on $f_1(n)$ which seems to tight.

For every $1 \leq r < n$ the graph $G = K_r + \overline{K_{n-r}}$ (see, e.g. [6]) satisfies

$$\mu_1(G) + \mu_1(\overline{G}) = \frac{r-1}{2} + \sqrt{nr - \frac{3r^2 + 2r - 1}{4}} + n - r - 1 = n - \frac{r+3}{2} + \sqrt{nr - \frac{3r^2 + 2r - 1}{4}}.$$

The right-hand side of this inequality is increasing in r for $0 \leq r \leq (n-1)/3$ and we find that

$$f_1(n) > \frac{4n}{3} - 2.$$

This gives some evidence for the following conjecture.

Conjecture 4

$$f_1(n) = \frac{4n}{3} + O(1).$$

We conclude this section with an improvement of the lower bound in (1). Using the first of inequalities (6) we obtain

$$\begin{aligned} \mu_1(G) + \mu_1(\overline{G}) &\geq n - 1 + \frac{s^2(G)}{2n^2} \left(\frac{1}{\sqrt{2e(G)}} + \frac{1}{\sqrt{2e(\overline{G})}} \right) \geq \\ &\geq n - 1 + \sqrt{2} \frac{s^2(G)}{n^3}. \end{aligned}$$

3 A class of graphs

In this section we shall describe a class of graphs that give the right order of $f_2(G)$ and, we believe, also of $f_n(G)$.

Let $n \geq 4$. Partition $[n]$ in 4 classes A, B, C, D so that $|A| \geq |B| \geq |C| \geq |D| \geq |A| - 1$. Join every two vertices inside A and D , join each vertex in B to each vertex in $A \cup C$, join each vertex in D to each vertex in C . Write $G(n)$ for the resulting graph.

Note that if n is divisible by 4, the sets A, B, C, D have equal cardinality and we see that $G(n)$ is isomorphic to its complement.

Our main goal to the end of this section is to estimate the eigenvalues of $G(n)$. Write $ch(A)$ for the characteristic polynomial of a matrix A . The following general theorem holds.

Theorem 5 *Suppose G is a graph and $V(G) = \cup_{i=1}^k V_i$ is a partition in sets of size n such that*

- (i) *for all $1 \leq i \leq k$, either $e(V_i) = \binom{n}{2}$ or $e(V_i) = 0$;*
- (ii) *for all $1 \leq i < j \leq k$, either $e(V_i, V_j) = n^2$ or $e(V_i, V_j) = 0$.*

Let the sets $V_1, \dots, V_p$ be independent and $V_{p+1}, \dots, V_k$ induce a complete graph. Then for the characteristic polynomial of the adjacency matrix of G we have

$$ch(A(G)) = x^{pn-p} (1-x)^{(k-p)n-(k-p)} ch(R),$$

where $R = (r_{ij})$ is a $k \times k$ matrix such that

$$r_{ij} = \begin{cases} 0 & \text{if } i \neq j \text{ and } e(V_i, V_j) = 0 \\ n & \text{if } i \neq j \text{ and } e(V_i, V_j) = n^2 \\ 0 & \text{if } i = j \text{ and } e(V_i) = 0 \\ n-1 & \text{if } i = j \text{ and } e(V_i) = \binom{n}{2} \end{cases}.$$

The proof of this theorem is a straight exercise in determinants, so we shall omit it.

If n is divisible by 4, say $n = 4k$, by Theorem 5, for the characteristic polynomial of $A(G(n))$ we have

$$ch(A(G(n))) = x^{2k-2} (1-x)^{2k-2} \begin{bmatrix} k-1-x & k & 0 & 0 \\ k & -x & k & 0 \\ 0 & k & -x & k \\ 0 & 0 & k & k-1-x \end{bmatrix}.$$

By straightforward calculations, setting $a = 1 - 1/k$ and $y = x/k$, we see that

$$\begin{aligned} ch(A(G(n))) &= x^{2k-2} (1-x)^{2k-2} [(a-y)(y^2(a-y) + 2y - a) - (y^2 - ay - 1)] \\ &= x^{2k-2} (1-x)^{2k-2} (y^2 - (1+a)y - (1-a))(y^2 + (1-a)y - (a+1)). \end{aligned}$$

Hence, we find that

$$\begin{aligned} \mu_2(G) &= -\frac{1}{2} + \sqrt{\frac{1}{4} + 2 \left\lfloor \frac{n}{4} \right\rfloor^2 - \left\lfloor \frac{n}{4} \right\rfloor} \\ \mu_n(G) &= -\frac{1}{2} - \sqrt{\frac{1}{4} + 2 \left\lfloor \frac{n}{4} \right\rfloor^2 - \left\lfloor \frac{n}{4} \right\rfloor}. \end{aligned}$$

If n is not divisible by 4, we will give some tight estimates of $\mu_2(G)$ and $\mu_n(G)$. Notice first that $G(4 \lfloor n/4 \rfloor)$ is an induced graph of $G(n)$ which in turn is an induced graph of $G(4 \lceil n/4 \rceil)$. Thus the adjacency matrix of $G(4 \lfloor n/4 \rfloor)$ is a principal submatrix of the adjacency matrix of $G(n)$ which in turn is a principal submatrix of the adjacency matrix of $G(4 \lceil n/4 \rceil)$. Since the eigenvalues of a matrix and its principal matrices are interlaced ([3], Theorem 4.3.15), we obtain

$$-\frac{1}{2} + \sqrt{\frac{1}{4} + 2 \left\lfloor \frac{n}{4} \right\rfloor^2 - \left\lfloor \frac{n}{4} \right\rfloor} \leq \mu_2(G) \leq -\frac{1}{2} + \sqrt{\frac{1}{4} + 2 \left\lceil \frac{n}{4} \right\rceil^2 - \left\lceil \frac{n}{4} \right\rceil}, \quad (10)$$

$$-\frac{1}{2} - \sqrt{\frac{1}{4} + 2 \left\lfloor \frac{n}{4} \right\rfloor^2 - \left\lfloor \frac{n}{4} \right\rfloor} \leq \mu_n(G) \leq -\frac{1}{2} - \sqrt{\frac{1}{4} + 2 \left\lceil \frac{n}{4} \right\rceil^2 - \left\lceil \frac{n}{4} \right\rceil}. \quad (11)$$

4 The asymptotics of $f_2(n)$

In this section we shall prove inequalities (4). From (10) we readily have

$$f_2(n) \geq -\frac{1}{2} + \sqrt{\frac{1}{4} + 2 \left\lfloor \frac{n}{4} \right\rfloor^2 - \left\lfloor \frac{n}{4} \right\rfloor} > \frac{\sqrt{2}}{2}n - 3,$$

so all we need to prove is that $f_2(n) \leq n/\sqrt{2}$.

By (8) we have

$$\mu_1^2(G) + \mu_2^2(G) + \mu_n^2(G) + \mu_1^2(\overline{G}) + \mu_2^2(\overline{G}) + \mu_n^2(\overline{G}) \leq n(n-1). \quad (12)$$

By Weyl's inequalities ([3], p. 181), for every graph G of order n , we have

$$\mu_2(G) + \mu_n(\overline{G}) \leq \mu_2(K_n) = -1.$$

Hence, using $\mu_2 \geq 0$ and $\mu_n \leq -1$ we obtain

$$\mu_2^2(G) \leq \mu_n^2(\overline{G}) + 2\mu_n(\overline{G}) + 1 < \mu_n^2(\overline{G}).$$

Hence, from (12) and $\mu_1(G) + \mu_1(\overline{G}) \geq n-1$, we find that

$$\frac{(n-1)^2}{2} + 2\mu_2^2(G) + 2\mu_2^2(\overline{G}) \leq \mu_1^2(G) + \mu_2^2(G) + \mu_n^2(G) + \mu_1^2(\overline{G}) + \mu_2^2(\overline{G}) + \mu_n^2(\overline{G}) \leq n(n-1).$$

After some algebra, we deduce that

$$\mu_2(G) + \mu_2(\overline{G}) \leq \frac{\sqrt{2}}{2}n,$$

completing the proof of inequalities (4).

5 Bounds on $f_n(n)$

In this section we shall prove inequalities (5). From (11), as above, we have

$$f_n(n) > \frac{\sqrt{2}}{2}n - 3.$$

We believe that, in fact, the following conjecture is true.

Conjecture 6

$$f_n(G) = \frac{\sqrt{2}n}{2} + O(1).$$

However we can only prove that $f_n(G) < (\sqrt{3}/2)n$ which is implied by the following theorem.

Theorem 7 *For every graph G of order n ,*

$$\mu_n^2(G) + \mu_n^2(\overline{G}) \leq \frac{3}{8}n^2.$$

Proof Indeed, suppose $(u_1, \dots, u_n)$ and $(w_1, \dots, w_n)$ are eigenvectors to $\mu_n(G)$ and $\mu_n(\overline{G})$. Let

$$U = \{i : u_i > 0\}, \quad W = \{i : w_i > 0\}.$$

Setting $V = [n]$, we clearly have $\mu_n^2(G) \leq E_G(U, V \setminus U)$ and $\mu_n^2(\overline{G}) \leq E_{\overline{G}}(W, V \setminus W)$. Since $E_G(U, V \setminus U) \cap E_{\overline{G}}(W, V \setminus W) = \emptyset$, we see that the graph

$$G' = (V, E_G(U, V \setminus U) \cup E_{\overline{G}}(W, V \setminus W))$$

is at most 4-colorable and hence G' contains no 4-cliques. By Turán's theorem (e.g., see [1]), we obtain $e(G') \leq (3/8)n^2$, completing the proof. $\square$

6 Bounds on $f_k(n)$, $2 < k < n$

In this section we shall give simple bounds on $f_k(n)$ for $2 < k < n$. Write for the Turán graph of order n with k classes. Recall that $T_k(n)$ is a complete k -partite graph whose vertex classes differ by at most 1 in size. We assume that k is fixed and n is large enough. Since $\mu_k(T_k(n)) = 0$ and $\mu_{n-k}(T_k(n)) \leq -\lfloor n/k \rfloor$ for n large, we immediately have

$$\begin{aligned} f_k(n) &\geq \lfloor n/k \rfloor - 1, \\ f_{n-k}(n) &\geq \lfloor n/k \rfloor + 1. \end{aligned}$$

We next turn to upper bounds on $f_k(n)$.

Theorem 8 *For any fixed k and any graph G of sufficiently large order n ,*

$$|\mu_k(G)| + |\mu_k(\overline{G})| < \sqrt{\frac{2}{k}}n. \quad (13)$$

and

$$|\mu_{n-k}(G)| + |\mu_{n-k}(\overline{G})| < \sqrt{\frac{2}{k}}n. \quad (14)$$

Proof Set $e(G) = m$. Our first goal is to prove that $|\mu_k(G)| \leq \sqrt{2e(G)/k}$. If $\mu_k(G) \geq 0$, we have in view of (8)

$$k\mu_k^2(G) \leq \sum_{i=1}^n \mu_i^2(G) = 2m.$$

If $\mu_k(G) < 0$ and $|\mu_k(G)| > \sqrt{2m/k}$ then

$$\sum_{i=1}^n \mu_i^2(G) \geq (n-k)\mu_k^2(G) > 2m \frac{n-k}{k} > 2m,$$

a contradiction. Hence, $|\mu_k(G)| \leq \sqrt{2e(G)/k}$, and, by symmetry, $|\mu_k(\overline{G})| \leq \sqrt{2e(\overline{G})/k}$.
Now

$$|\mu_k(G)| + |\mu_k(\overline{G})| \leq \sqrt{2e(G)/k} + \sqrt{2e(\overline{G})/k} \leq \sqrt{\frac{2}{k}n(n-1)} < \sqrt{\frac{2}{k}n},$$

proving inequality (13). The proof of inequality (14) goes along the same lines, so we will omit it. $\square$

References

- [1] B. Bollobás, *Modern Graph Theory*, Graduate Texts in Mathematics, 184, Springer-Verlag, New York (1998), xiv+394 pp.
- [2] D. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1980, 368 pp.
- [3] R. Horn and C. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985. xiii+561 pp.
- [4] Y. Hong, On the spectral radius and the genus of graphs, J. Combin. Theory Ser. B 65 (2) (1995), 262-268.
- [5] Y. Hong, J. Shu, A sharp upper bound for the spectral radius of the Nordhaus-Gaddum type, Discr. Math. 211 (2000), 229-232.
- [6] Y. Hong, J. Shu, and K. Fang, A sharp upper bound of the spectral radius of Graphs, J. Combin. Theory Ser. B 81 (2001), 177-183.
- [7] X. Li, The relations between the spectral radius of the graphs and their complements, J. North China Technol. Inst. 17 (4) (1996), 297-299.
- [8] V. Nikiforov, Some inequalities for the largest eigenvalue of a graph. Combin. Probab. Comput. 11 (2002), 179-189.
- [9] V. Nikiforov, Eigenvalues and degree deviation, submitted.
- [10] E. Nosal, Eigenvalues of Graphs, Master's thesis, University of Calgary, 1970.
- [11] B. Zhou, A note about the relations between the spectral radius of graphs and their complements, Pure Appl. Math. 13 (1) (1997), 15-18.